



PubMed Nucleotide Protein Genome Structure PopSet Taxonomy OMIM

Search PubMed

for

Limits Preview/Index History Clipboard

Abstract

About Entrez

Entrez PubMed  
Overview  
Help | FAQ  
New/Noteworthy

PubMed Services  
Journal Browser  
MeSH Browser  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
Cubby NEW

Related Resources  
Order Documents  
Grateful Med  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov

Privacy Policy

1: *J Appl Toxicol* 1997 Sep-Oct;17(5):337-43

Related Articles, Books, LinkOut

**InterScience**

**Cadmium and selenium in blood and urine related to smoking habits and previous exposure to mercury vapour.**

**Ellingsen DG, Thomassen Y, Aaseth J, Alexander J**

Department of Occupational and Environmental Medicine, Telemark Central Hospital, Skien, Norway.

The object of this work was to investigate possible interactions of mercury, cadmium and selenium in humans. Selenium and cadmium in blood and urine were determined in this cross-sectional study of 130 males, of whom 77 had been previously exposed to mercury vapour at a chloralkali plant. Of the participants, 61.5% were smokers and 16.2% were never-smokers. The concentration of selenium in blood (B-Se) was significantly lower in subjects currently smoking more than 50 g of tobacco per week compared to never-smokers, whereas the concentration of cadmium in blood (B-Cd) was significantly higher in all categories of current smokers. In the multiple linear regression analysis, B-Se as a dependent variable was negatively associated with B-Cd, whereas current smoking habits were not included in the model as a predictor variable. In contrast, B-Cd as a dependent variable was positively associated with current as well as previous smoking habits, and negatively with both B-Se and the 'cumulative dose' of previous mercury vapour exposure. The concentration of selenium in blood was also negatively associated with B-Cd in the group of never-smokers (Spearman's  $r = -0.80$ ;  $P < 0.001$ ). In conclusion, these results suggest a depressive effect of cadmium on the concentration of selenium in blood, while smoking alone did not operate as a true predictor for this effect. Furthermore, previous exposure to mercury apparently modifies the concentration of cadmium in blood.

PMID: 9339747, UI: 97479666

Abstract

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services  
Freedom of Information Act | Disclaimer